




Golden Pro™ 200 EC

Golden Pro™ 200EC is a broad spectrum insecticide for the control of sucking and chewing insects on a wide range of crops



Active Ingredients


Profenofos 40% + Cypermethrin 4% Emulsifiable Concentrate



For Use on a wide range of Crops

Maize, Wheat, Cotton, Soybeans, Beans, Peas, Groundnuts, Potatoes, Tomatoes, Peppers and Onions


Cypermethrin



- Pyrethroid insecticide, IRAC Group 3A

A synthetic pyrethroid insecticide with fast action and knock down effect. It has both contact and stomach action activity.

Profenofos



- Organophosphate insecticide, IRAC Group 1B

An Organophosphate insecticide with ovicidal and acaricidal activity. It has both contact and stomach action with additional translaminar properties.

Golden Pro™ 200EC characteristics

- Long lasting effect which is sped up by high temperatures and high relative humidity
- Does not affect the flavour and odour of the treated crops
- Effective against a wide range of differentiating insects (aphids, thrips, caterpillars, beetles, etc)
- Quick knock-down effect
- Synergistic power with both contact and stomach insecticide action
- Combination of insecticides with two different mode of actions
- Translaminar activity, leads to increased efficiency against insects in "hidden" spots and the lower surface of the leaves



Application Rate

Crop	Pest	Application Rate	Notes
Cereal (Maize, Wheat)	Lepidoptera pests, Aphids, Thrips	400 – 600 ml/ha ; 50 – 80 ml/100 l water	Apply upon pest appearance. Ensure thorough coverage of the pest canopy.
Cotton	Lepidoptera pests incl. Bollworms, Aphids, Spidermites, Thrips		
Soybeans, Beans, Peas, Groundnuts	Aphids, Lepidoptera pests, Thrips, Whiteflies		
Vegetables (Potatoes, Tomatoes, Peppers, Onions)	Aphids, Lepidoptera pests, Thrips, Whiteflies, Spidermites		

- In order to delay insecticide resistance, always alternate with products with different mode of action.
- Mix the required quantity of **Golden Pro™ 200EC** with water in a spray tank while stirring or with agitator in motion
- Apply when the insect infestation reaches economic threshold level
- Repeat application at 10 - 15 days interval if infestation persists.



Disclaimer: This product must be used strictly as directed and in accordance with the instructions appearing on the label. The supplier accepts no liability other than as indicated in the Supplier's 'terms and conditions of sale' and the warranty clauses indicated on the product label. Sellers guarantee is limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

KENAGRO
KENAGRO LTD,
 Agias Fylaxeos & Zinonos Rossidi 2,
 2/F, P.C. 3082 Limassol, Cyprus
 E-mail: info@kenagroLtd.com